

Gender Affirming Surgery

Policy MP-002

Origination Date: 04/24/2018

Reviewed/Revised Date: 08/23/2021

Next Review Date: 08/23/2022

Current Effective Date: 10/23/2021

Disclaimer:

1. Policies are subject to change in accordance with State and Federal notice requirements.
2. Policies outline coverage determinations for U of U Health Plans Commercial, and Healthy U (Medicaid) plans. Refer to the "Policy" section for more information.

Description:

Gender affirming surgery is part of the spectrum of care considered for individuals with gender dysphoria, a condition in which a person feels a strong and persistent identification with the opposite gender accompanied with a severe sense of discomfort in their own gender. People with gender dysphoria often report a feeling of being born the wrong gender they physically appear to be. Gender affirming surgery is not a single procedure, but part of a complex process involving multiple medical, psychiatric, and surgical modalities working in conjunction with each other and the patient to achieve successful behavioral and medical outcomes.

Policy Statement and Criteria

1. Commercial Plans

Criteria is divided by Breast/Chest Surgery and Genital Surgery

U of U Health Plans covers gender affirming surgery when all of the following criteria are met:

Criteria for Breast/Chest Surgery

- A. Patient is 18 years or older;
- B. The requested procedure is being performed by qualified physicians at an approved Center of Excellence with experience in the following services;
- C. The patient has been diagnosed with persistent and documented Gender Dysphoria
- D. Documentation that no significant medical or mental health concerns are present. If concerns are present, then documentation that they are reasonably controlled;
- E. Capacity to make a fully informed decision and to give consent for treatment;

- F. One letter provided by treating provider documenting the following:
- i. The patient's general identifying characteristics; and
 - ii. The initial and evolving gender, sexual, and other psychiatric diagnoses; and
 - iii. The duration of the patient's professional relationship with the provider or previous providers including the type of psychotherapy or evaluation that the patient underwent; and
 - iv. The eligibility criteria that have been met and the behavioral health professional's rationale for surgery; and
 - v. The degree to which the patient has followed the eligibility criteria to date and the likelihood of future compliance; and
 - vi. Whether the author of the report is part of a gender team.

Criteria for Genital Surgery

Hysterectomy and Salpingo-Oophorectomy in FtM Patients and Orchiectomy in MtF Patients:

- A. Patient is 18 years or older;
- B. The requested procedure is being performed by qualified physicians at an approved Center of Excellence with experience in the following services;
- C. The patient has been diagnosed with persistent and well documented Gender Dysphoria;
- D. Documentation that significant medical or mental health concerns, if present, are well controlled;
- E. For these patients without a medical contraindication, the patient has undergone a minimum of 12 months of continuous hormonal therapy;
- F. The patient has completed a minimum of 12 months of successfully living in the gender role that is congruent with their gender identity;
- G. Two letters provided by treating providers:
 - i. The first letter from the patient's physician or behavioral health provider, documenting the following:
 - a) The patient's general identifying characteristics; and
 - b) The initial and evolving gender, sexual, and other psychiatric diagnoses; and
 - c) The duration of the patient's professional relationship with the provider or previous providers including the type of psychotherapy or evaluation that the patient underwent; and

- d) The eligibility criteria that have been met and the behavioral health professional's rationale for surgery; and
 - e) The degree to which the patient has followed the eligibility criteria to date and the likelihood of future compliance; and
 - f) Whether the author of the report is part of a gender team.
- ii. A second letter from a different physician or behavioral health provider familiar with the patient's treatment and the psychological aspects of Gender Dysphoria, corroborating the information provided in the first letter;
 - iii. When one of the signatories on the letters indicated above is not the treating surgeon, a letter from the surgeon confirming that they have personally communicated with the treating therapist and or physician, as well as the patient, and confirming that the patient meets the above criteria, understands the ramifications and possible complications of surgery, and that the surgeon feels that the patient is likely to benefit from surgery.

Metoidioplasty or Phalloplasty in FtM Patients and Vaginoplasty in MtF Patients:

- A. Patient is 18 years or older;
- B. The requested procedure is being performed by qualified physicians at an approved Center of Excellence with experience in the following services;
- C. The patient has been diagnosed with persistent and well documented Gender Dysphoria;
- D. For these patients without a medical contraindication, the patient has undergone a minimum of 12 months of continuous hormonal therapy;
- E. Two letters provided by treating providers:
 - i. The first letter from the patient's physician or behavioral health provider, documenting the following:
 - a) The patient's general identifying characteristics; and
 - b) The initial and evolving gender, sexual, and other psychiatric diagnoses; and
 - c) The duration of the patient's professional relationship with the provider or previous providers including the type of psychotherapy or evaluation that the patient underwent; and
 - d) The eligibility criteria that have been met and the behavioral health professional's rationale for surgery; and
 - e) The degree to which the patient has followed the eligibility criteria to date and the likelihood of future compliance; and

- f) Whether the author of the report is part of a gender team.
- ii. A second letter from a different physician or behavioral health provider familiar with the patient's treatment and the psychological aspects of Gender Dysphoria, corroborating the information provided in the first letter;
 - iii. When one of the signatories on the letters indicated above is not the treating surgeon, a letter from the surgeon confirming that they have personally communicated with the treating therapist and or physician, as well as the patient, and confirming that the patient meets the above criteria, understands the ramifications and possible complications of surgery, and that the surgeon feels that the patient is likely to benefit from surgery.

Gender affirming surgery may include any of the following procedures:

Male-to-Female Procedures:

- Breast implantation/augmentation
- Orchiectomy
- Penectomy
- Vaginoplasty and associate hair removal if necessary for external skin transpositioned into area
- Clitoroplasty
- Labiaplasty

Female-to- Male Procedures:

- Subcutaneous mastectomy without nipple rearrangement surgery
- Hysterectomy
- Salpingo-oophorectomy
- Vaginectomy
- Metoidioplasty
- Scrotoplasty
- Urethroplasty
- Placement of testicular prostheses Phalloplasty

In the case of non-binary individuals, above surgeries will be considered to align their gender preference.

Gender affirming surgery is **not covered** and considered cosmetic when used to improve the gender specific appearance of a patient who has undergone or is planning to undergo gender affirming surgery.

The following surgeries are considered cosmetic (may not be all inclusive):

- 1) Abdominoplasty
- 2) Blepharoplasty
- 3) Brow Lift
- 4) Cheek/Malar Implants
- 5) Chin/Nose Implants
- 6) Collagen Injections
- 7) Facial bone reconstruction
- 8) Face lift
- 9) Forehead Lift
- 10) Calf Lift
- 11) Hair removal/hairplasty including medications that cause hair loss or growth unless area of hair removal is skin taken from external source used for creation of artificial vagina
- 12) Hair Transplantation
- 13) Lip Reduction
- 14) Liposuction
- 15) Mastopexy
- 16) Neck Tightening
- 17) Pectoral Implants
- 18) Reduction thyroid chondroplasty
- 19) Rhinoplasty
- 20) Voice modification surgery
- 21) Voice Therapy/Lessons

It is important to note this policy **DOES NOT** apply to individuals with congenital deformities/anomalies or genetic abnormalities resulting in genitalia requiring correction.

2. Medicaid Plans

Coverage is determined by the State of Utah Medicaid program; if Utah State Medicaid has no published coverage position and InterQual criteria are not available, the U of U Health Plans Commercial criteria will apply. For the most up-to-date Medicaid policies and coverage, please visit their website at

<http://health.utah.gov/medicaid/manuals/directory.php> or the [Utah Medicaid code Look-Up tool](#)

3. Medicare Plans

Coverage is determined by the Centers for Medicare and Medicaid Services (CMS); if a coverage determination has not been adopted by CMS and InterQual criteria are not available, the U of U Health Plans Commercial criteria will apply. For the most up-to-date Medicare policies and coverage, please visit their search website at:

<http://www.cms.gov/medicare-coverage-database/overview-and-quick-search.aspx?from2=search1.asp&> or [the manual website](#)

Clinical Rationale

Gender Dysphoria, previously referred to as Gender Identity Disorder or transsexualism, is a condition wherein an individual's experienced/expressed gender is incongruent with the assigned gender and causes clinically significant distress and impairment. Gender dysphoria should not be confused with cross dressing (Transvestitism), refusal to accept non-heterosexual orientation, psychotic delusions or personality disorders.

Gender-affirming surgical treatment differs depending upon the assigned sex at birth of the patient and each patient's needs to diminish gender dysphoria. Not every patient wants every type of gender-affirming surgery. Male-to-female patients, also known as "transwomen," may seek genital surgery (removal of the testicles only; removal of testicles and penis and the creation of pseudo vagina, clitoris, and labia) or facial feminization surgery (tracheal shave). Female-to-male patients, known as "transmen," may seek genital surgery to remove the uterus, ovaries, and vagina, creation of a neophallus, and scrotum with scrotal prostheses) and "top surgery", also called chest masculinization surgery with bilateral mastectomy.

The guideline criteria above are based upon; 1) the Diagnostic and Statistical manual of Mental Disorders, Fifth Edition, (DSM-V, 2013) criteria for the diagnosis of Gender Dysphoria; and 2) the Standards of Care (SOC) for Transgender and Gender Non-Conforming People, Version 7, published by the World Professional Association for Transgender Health (2012). Both of these references are widely accepted as definitive documents in the area of Gender Dysphoria treatment and cited in numerous articles by respected authors. The SOC criteria have been adopted in several countries as the standard of care for the treatment of Gender Dysphoria, including hormone therapy and gender affirming surgery.

The criteria of the SOC are supported by evidence-based peer-reviewed journal publications. Several studies have shown that extensive long-term trials of hormonal therapy and real-life experiences, as well as social support and acceptance by peer and family groups, greatly improve psychological and social outcomes in patients undergoing gender affirming surgery (Eldh, 1997; Landen, 1998). A study reported by Monstrey and colleagues (2001) described the importance of close cooperation between the many medical and behavioral specialties required for proper treatment of patients with Gender Dysphoria who wish to undergo gender affirming surgery. Similar findings were reported earlier by Schlatterer et al. in 1996. One study of 188 patients undergoing gender affirming surgery found that dissatisfaction with surgery was highly associated with sexual preference, psychological co-morbidity, and poor preoperative body image and satisfaction (Smith, 2005).

Undertaking gender affirming surgery is obviously a very serious decision. The procedures present significant medical and psychological risks, and results are irreversible. A step-wise approach to therapy for Gender Dysphoria, including accurate diagnosis and long-term treatment by multidisciplinary team including behavioral, medical and surgical specialists, has been shown to provide the best results. As

with any treatment including psychiatric disorders, a thorough behavioral analysis by a qualified practitioner is needed. Once a diagnosis of Gender Dysphoria is established, treatment with hormone therapy and establishment of real-life transgender experience may be warranted. Gender affirming surgery should be considered only after such trials have been undertaken, evaluated and confirmed. Hormone therapy should be administered under on-going medical supervision and is important in beginning the gender transition process by altering body hair, breast size, skin appearance and texture, body fat distribution, and the size and function of sex organs. Additionally, real-life experience is important to validate the patient's desire and ability to incorporate into their desired gender role within their social network and daily environment. This generally involves gender-specific appearance (clothing, hairstyle, etc.), involvement in various activities in the desired gender role including work or academic settings, legal acquisition of a gender appropriate first name, and acknowledgement by others of their new gender role.

Once these treatments steps have been established and stable for at least 12 months, a patient may be considered for gender affirming surgery.

For both transmen and transwomen, additional surgeries have been proposed to improve the gender appropriate appearance of the patient. Procedures such as breast augmentation, liposuction, Adam's apple reduction (tracheal shave), rhinoplasty, facial reconstruction, and others have no medically necessary role in gender identification and are considered cosmetic in nature.

The World Professional Association for Transgender Health (WPATH) is a multidisciplinary professional society representing the specialties of medicine, psychology, social sciences and law that has published clinical guidelines regarding health services for patients with gender disorders. In 2012, WPATH updated their evidence and consensus-based guideline regarding, the Standards of Care (SOC) for the Health of Transsexual, Transgender, and Gender Nonconforming Peoples.

WPATH guidelines indicate that surgical treatments can be initiated by a referral from a qualified mental health professional. One or two referrals may be required depending upon the type of surgery requested. "The mental health professional provides documentation—in the chart and/or referral letter—of the patient's personal and treatment history, progress, and eligibility." WPATH guidelines specifically recommend the following:

- One referral from a qualified mental health professional is needed for breast/chest surgery (e.g., mastectomy, chest reconstruction, or augmentation mammoplasty).
- Two referrals—from qualified mental health professionals who have independently assessed the patient—are needed for genital surgery (i.e., hysterectomy/salpingo-oophorectomy, orchiectomy, genital reconstructive surgeries).

WPATH lists the following criteria for mastectomy and creation of a male chest in FTM patients:

1. Persistent, well-documented gender dysphoria;
2. Capacity to make a fully informed decision and to consent for treatment;
3. Age of majority in a given country (or with 2-parent consent);
4. If significant medical or mental health concerns are present, they must be reasonably well controlled.

Hormone therapy is not a prerequisite.

WPATH lists the following criteria for genital surgery:

1. Persistent, well-documented gender dysphoria;

2. Capacity to make a fully informed decision and to consent for treatment;
3. Age of majority in a given country;
4. If significant medical or mental health concerns are present, they must be reasonably well controlled; and
5. 12 continuous months of hormone therapy as appropriate to the patient's gender goals (unless hormones are not clinically indicated for the individual).

In addition, WPATH made specific recommendations regarding breast augmentation procedures:

The WPATH guideline recommends MTF patients undergo feminizing hormone therapy for a minimum of 12 months prior to augmentation surgery and lists specific criteria for breast augmentation (implants/lipofilling).

However, the classification of breast augmentation as a cosmetic versus reconstructive procedure has remained controversial. WPATH guidelines note that although breast appearance may be considered an important secondary sex characteristic, opinions diverge regarding whether augmentation is considered cosmetic or reconstructive. In addition, WPATH indicates that, "breast presence or size is not involved in the legal definitions of sex and gender and is not necessary for reproduction."

In 2009, the Endocrine Society in conjunction with European Society of Endocrinology, European Society for Pediatric Endocrinology, Lawson Wilkins Pediatric Endocrine Society, and World Professional Association, published the only evidence-based guidelines regarding the treatment of transsexual persons. The guideline employed transparent methods for evidence review and for rating the quality of evidence. All recommendations were based upon evidence which was rated to be low quality. These guidelines were updated in 2017. The consortium made the following recommendations:

- 1) We recommend that a patient pursue genital gender affirming surgery only after the MHP and the clinician responsible for endocrine transition therapy both agree that surgery is medically necessary and would benefit the patient's overall health and/or well-being.
- 2) We advise that clinicians approve genital genderaffirming surgery only after completion of at least 1 year of consistent and compliant hormone treatment, unless hormone therapy is not desired or medically contraindicated.
- 3) We advise that the clinician responsible for endocrine treatment and the primary care provider ensure appropriate medical clearance of transgender individuals for genital gender-affirming surgery and collaborate with the surgeon regarding hormone use during and after surgery.
- 4) We recommend that clinicians refer hormone treated transgender individuals for genital surgery when:
 - (a) the individual has had a satisfactory social role change, (b) the individual is satisfied about the hormonal effects, and (c) the individual desires definitive surgical changes.
- 5) We suggest that clinicians delay gender-affirming genital surgery involving gonadectomy and/or hysterectomy until the patient is at least 18 years old or legal age of majority in his or her country.
- 6) We suggest that clinicians determine the timing of breast surgery for transgender males based upon the physical and mental health status of the individual.
- 7) There is insufficient evidence to recommend a specific age requirement.

Coding for Breast and Nipple Reconstruction. The CPT codes for mastectomy (CPT codes 19303 and 19304) are for breast cancer, and are not appropriate to bill for reduction mammoplasty for female to male (transmasculine) gender affirmation surgery. CPT 2020 states that “Mastectomy procedures (with the exception of gynecomastia [19300]) are performed either for treatment or prevention of breast cancer” and that “Code 19303 describes total removal of ipsilateral breast tissue with or without removal of skin and/or nipples (eg, nipple-sparing), for treatment or prevention of breast cancer.” There are important differences between a mastectomy for breast cancer and a mastectomy for gender affirmation. The former requires careful attention to removal of all breast tissue to reduce the risk of cancer. By contrast, careful removal of all breast tissue is not essential in mastectomy for gender affirmation. In mastectomy for gender affirmation, the nipple areola complex typically can be preserved.

Some have tried to justify routinely billing CPT code 19350 for nipple reconstruction at the time of mastectomy for gender affirming based upon the frequent need to reduce the size of the areola to give it a male appearance. However, the nipple reconstruction as defined by CPT code 19350 describes a much more involved procedure than areola reduction. The typical patient vignette for CPT code 19350, according to the AMA, is as follows: “The patient is measured in the standing position to ensure even balanced position for a location of the nipple and areola graft on the right breast. Under local anesthesia, a Skate flap is elevated at the site selected for the nipple reconstruction and constructed. A full-thickness skin graft is taken from the right groin to reconstruct the areola. The right groin donor site is closed primarily in layers.”

The AMA vignette for CPT code 19318 (breast reduction) clarifies that this CPT code includes the work that is necessary to reposition and reshape the nipple to create an aesthetically pleasing result, as is necessary in female to male breast reduction. “The physician reduces the size of the breast, removing wedges of skin and breast tissue from a female patient. The physician makes a circular skin incision above the nipple, in the position to which the nipple will be elevated. Another skin incision is made around the circumference of the nipple. Two incisions are made from the circular cut above the nipple to the fold beneath the breast, one on either side of the nipple, creating a keyhole shaped skin and breast incision. Wedges of skin and breast tissue are removed until the desired size is achieved. Bleeding vessels may be ligated or cauterized. The physician elevates the nipple and its pedicle of subcutaneous tissue to its new position and sutures the nipple pedicle with layered closure. The remaining incision is “repaired with layered closure” (EncoderPro, 2021). CPT code 19350 does not describe the work that is being done, because that code describes the actual construction of a new nipple.

Therefore, U of U Health Plans considers nipple reconstruction, as defined by CPT code 19350, as cosmetic and/or not medically necessary for mastectomy for transmasculine gender affirming, and that CPT code 19318 includes the extra work that may be necessary to reshape the nipple and create an aesthetically pleasing male chest.

Applicable Coding

Covered: For the conditions outlined above for plans with gender affirming supplemental coverage.

CPT Codes

00402 Anesthesia for procedures on the integumentary system on the extremities, anterior trunk and perineum; reconstructive procedures on breast (eg, reduction or augmentation mammoplasty, muscle flaps)

00926 Anesthesia for procedures on male genitalia (including open urethral procedures); radical orchiectomy, inguinal

11980 Subcutaneous hormone pellet implantation (implantation of estradiol and/or testosterone pellets beneath the skin)

15771 Grafting of autologous fat harvested by liposuction technique to trunk, breasts, scalp, arms, and/or legs; 50 cc or less injectate

15772 Grafting of autologous fat harvested by liposuction technique to trunk, breasts, scalp, arms, and/or legs; each additional 50 cc injectate, or part thereof (List separately in addition to code for primary procedure)

19318 Breast reduction

19324 Mammoplasty, augmentation; without prosthetic implant *(Deleted 01/01/2021)*

19325 Mammoplasty, augmentation; with prosthetic implant

53430 Urethroplasty, reconstruction of female urethra

54125 Amputation of penis; complete

54400 Insertion of penile prosthesis; non-inflatable (semi-rigid)

54401 Insertion of penile prosthesis; inflatable (self-contained)

54520 Orchiectomy, simple (including subcapsular), with or without testicular prosthesis, scrotal or inguinal approach

54660 Insertion of testicular prosthesis (separate procedure)

54690 Laparoscopy, surgical; orchiectomy

55150 Resection of scrotum

55175 Scrotoplasty; simple

55180 Scrotoplasty; complicated

55899 Unlisted procedure, male genital system

55970 Intersex surgery; male to female

55980 Intersex surgery; female to male

56625 Vulvectomy simple; complete

56800 Plastic repair of introitus

56805 Clitoroplasty for intersex state

57110 Vaginectomy, complete removal of vaginal wall;

57291 Construction of artificial vagina; without graft

57292 Construction of artificial vagina; with graft

- 57335** Vaginoplasty for intersex state
- 58150** Total abdominal hysterectomy (corpus and cervix), with or without removal of tube(s), with or without removal of ovary(s);
- 58262** Vaginal hysterectomy, for uterus 250 g or less; with removal of tube(s), and/or ovary(s)
- 58291** Vaginal hysterectomy, for uterus greater than 250 g; with removal of tube(s) and/or ovary(s)
- 58552** Laparoscopy, surgical, with vaginal hysterectomy, for uterus 250 g or less; with removal of tube(s) and/or ovary(s)
- 58554** Laparoscopy, surgical, with vaginal hysterectomy, for uterus greater than 250 g; with removal of tube(s) and/or ovary(s)
- 58571** Laparoscopy, surgical, with total hysterectomy, for uterus 250 g or less; with removal of tube(s) and/or ovary(s)
- 58573** Laparoscopy, surgical, with total hysterectomy, for uterus greater than 250 g; with removal of tube(s) and/or ovary(s)
- 58661** Laparoscopy, surgical; with removal of adnexal structures (partial or total oophorectomy and/or salpingectomy)
- 58720** Salpingo-oophorectomy, complete or partial, unilateral or bilateral (separate procedure)
- 58940** Oophorectomy, partial or total, unilateral or bilateral;

(CPT Codes Not Covered for Gender Affirming Surgery)

- 19303** Mastectomy, simple, complete
- 19350** Nipple/areola reconstruction

HCPCS Codes

- C1789** Prosthesis, breast (implantable)
- C1813** Prosthesis, penile, inflatable
- L8600** Implantable breast prosthesis, silicone or equal
- S0189** Testosterone pellet, 75 mg

References:

1. Becker S, Bosinski H, Clement U, et al. German Standards for the Treatment and Diagnostic Assessment of Transsexuals. Issued by the German Society for Sexual Research, the Academy for Sexual Medicine and the Society for Sexual Sciences. *Int J Transgender* 1998; 2(4). Available at <http://www.symposion.com/jjt/jjtc0603.htm>. Accessed on November 6, 2006.
2. Bradley SJ, Zucker KJ. Gender identity disorder; A review of the past 10 years. *J Am Acad Child Adolesc Psychiatry*. 1997; 36(7):872-880.
3. Cohen-Kettenis PT, Gooren LJ. Transsexualism: a review of etiology, diagnosis and treatment. *J Psychosom Res*. 1999; 46(4):315-333. Centers for Medicare and Medicaid Services. National Coverage Determination: Transsexual, NCD #140.3.

Effective date not available. Available at: http://www.cms.hhs.gov/med/index_list.asp?list_type=pcd. Accessed on November 6, 2006.

4. Day P. Trans-gender reassignment surgery. New Zealand health technology assessment (NZHTA). The clearing house for health outcomes and Health technology assessment. February 2002; Volume 1 Number 1 Available at: http://nzhta.chmeds.ac.nz/publications/trans_gender.pdf. Assessed on November 6, 2006.
5. Eldh J, Berg A, Gustaffson M. Long-term follow up after sex reassignment surgery. *Scand J Plast Reconstr Surg Hand Surg*. 1997; 31(1):39-45.
6. Endocrine Treatment of Gender-Dysphoric/Gender-Incongruent Persons: An Endocrine Society Clinical Practice Guideline *The Journal of Clinical Endocrinology & Metabolism* ,Volume 102, Issue 11 1 November 2017 ,Pages 3869-3903
7. Endocrine Treatment of Transsexual Persons: An Endocrine Society Clinical Practice Guideline (2009)
8. Hayes, Inc. Hayes Medical Technology Directory. Sex reassignment surgery and associated therapies for the treatment of gender identity disorder. Lansdale, PA: Hayes, Inc.; December 2004. Search updated December 20, 2005.
9. Hepp U, Kraemer B, Schbyder U, miller N, Delsignore A. Psychiatric comorbidity in gender identity disorder. *J Psychosom Res*. 2005; 58(3): 259-261.
10. Kraus, C. (2015). "Classifying Intersex in DSM-5: Critical Reflections on Gender Dysphoria." *Arch Sex Behav* 44(5): 11471163.
11. Landen M, Walinder J, Hambert G, Lundstrom B. factors predictive of regret in sex reassignment. *Acta Psychiatr Scand*, 1998; 7(4):284-9.
12. Mate-Kole C, Freschi M, Robin A. A controlled study of psychological and social change after surgical fender reassignment in selected male transsexuals. *Br J Psychiatry*. 1990; 157:261-264.
13. Midence K, Hargreaves I. Psychosocial adjustment in male-to-female transsexuals: an overview of the research evidence. *J. Psychol*. 1997; 131(6):602-614.
14. Monstrey S, Hoebeke P, Dhont M, et al. Surgical therapy in transsexual patients: a multi-disciplinary approach. *Acta Chir Belg*. 2001; 101(5): 200-209.
15. Optum 360[®] EncoderPro.com (2021)
16. Parkes G, Hall I. Gender dysphoria and cross-dressing in people with intellectual disability: a literature review, *Ment Retard*. 2006; 44(4):260-71.
17. Schlatterer K, von Werder K, Stalla GK. Multistep treatment concept of transsexual patients. *Exp Clin Endocrinol Diabetes*. 1996; 104(6):413-419.
18. Selvaggi G, Ceulemans P, De Cuyper g, et al. gender identity disorder: general overview and surgical treatments for vaginoplasty in male-to-female transsexuals. *Plast Reconstr Surg*. 2005; 116(6):135e-145e.
19. Smith YL, Van Goozen SH, Kuiper AJ, Cohen-Kettenis PT. Sex Reassignment: outcomes and predictors of treatment for adolescent and adult transsexuals. *Psychol Med*. 2005; 35(1):89-99.
20. The Harry Benjamin International Gender Dysphoria Association. Standards of Care for Gender Identity Disorders. 6th Version. 2001 Feb. Available at: <http://www.hbigoa.org/Documents2/socv6.pdf>. Assessed on: November 6, 2006.
21. World Professional Association for Transgender Health (WPATH) V7 , 2012.

Disclaimer:

This document is for informational purposes only and should not be relied on in the diagnosis and care of individual patients. Medical and Coding/Reimbursement policies do not constitute medical advice, plan preauthorization, certification, an explanation of benefits, or a contract. Members should consult with appropriate health care providers to obtain needed medical advice, care, and treatment. Benefits and eligibility are determined before medical guidelines and payment guidelines are applied. Benefits are determined by the member's individual benefit plan that is in effect at the time services are rendered.

The codes for treatments and procedures applicable to this policy are included for informational purposes. Inclusion or exclusion of a procedure, diagnosis or device code(s) does not constitute or imply member coverage or provider reimbursement policy. Please refer to the member's contract benefits in effect at the time of service to determine coverage or non-coverage of these services as it applies to an individual member.

U of U Health Plans makes no representations and accepts no liability with respect to the content of any external information cited or relied upon in this policy. U of U Health Plans updates its Coverage Policies regularly, and reserves the right to amend these policies and give notice in accordance with State and Federal requirements.

No part of this publication may be reproduced, stored in a retrieval system or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from U of U Health Plans.

"University of Utah Health Plans" and its accompanying logo, and its accompanying marks are protected and registered trademarks of the provider of this Service and or University of Utah Health. Also, the content of this Service is proprietary and is protected by copyright. You may access the copyrighted content of this Service only for purposes set forth in these Conditions of Use.