

Drug Levels and Antibody Testing for Infliximab, Adalimumab, Vedolizumab and Ustekinumab

Policy MP-026

Origination Date: 07/15/2021

Reviewed/Revised Date: 07/15/2021

Next Review Date: 07/15/2022

Current Effective Date: 09/20/2021

Disclaimer:

1. Policies are subject to change in accordance with State and Federal notice requirements.
2. Policies outline coverage determinations for U of U Health Plans Commercial, and Healthy U (Medicaid) plans. Refer to the "Policy" section for more information.

Description:

Infliximab, adalimumab, ustekinumab, and vedolizumab are monoclonal antibody drugs used to treat inflammatory conditions, such as rheumatoid arthritis, psoriatic arthritis, juvenile idiopathic arthritis; inflammatory bowel disease (e.g., Crohn's disease, ulcerative colitis), ankylosing spondylitis, and plaque psoriasis. These biological agents are generally given to patients who fail conventional medical therapy, and are typically effective for the induction and maintenance of clinical remission. However, during use, some patients develop anti-drug antibodies (ADAs) that neutralize the anti-inflammatory action of these drugs. This response may diminish the potential long-term efficacy of these drugs.

The reasons for therapeutic failures remain a matter of debate but include accelerated drug clearance (pharmacokinetics) and neutralizing agent activity (pharmacodynamics) due to the development of ADAs. Another proposed mechanism to determine a patient's loss of response to the monoclonal antibody drugs is therapeutic drug monitoring (TDM). These blood tests are used to measure and monitor both the serum drug level and antibody level to these drugs. However, the detection and quantitative measurement of ADAs are difficult because of drug interference and identifying when antibodies have a neutralizing effect. Which leaves several obstacles to their regular use, including when to use TDM, how to accurately interpret and apply the results of such testing, and defining the optimal drug concentration thresholds and ranges to target.

Policy Statement and Criteria

1. Commercial Plans

U of U Health Plans does not cover testing for the measurement of antibodies and/or serum drug levels to monoclonal antibody (MAB) drugs, including anti-tumor necrosis factor (TNF) drugs, either alone or as a combination test, as it is considered investigational or unproven.

U of U Health Plans does NOT cover the following tests (may not be an all-inclusive list):

- Anser® ADA
- Anser® IFX
- Anser® VDZ
- Anser® UST
- Miraca Life Sciences' InformTx™
- DoseASSURE™-LabCorp
- LabCorp®-ECLIA

2. Medicaid Plans

Coverage is determined by the State of Utah Medicaid program; if Utah State Medicaid has no published coverage position and InterQual criteria are not available, the U of U Health Plans Commercial criteria will apply. For the most up-to-date Medicaid policies and coverage, please visit their website at:

<http://health.utah.gov/medicaid/manuals/directory.php> or the [Utah Medicaid code Look-Up tool](#)

3. Medicare Plans

Coverage is determined by the Centers for Medicare and Medicaid Services (CMS); if a coverage determination has not been adopted by CMS and InterQual criteria are not available, the U of U Health Plans Commercial criteria will apply. For the most up-to-date Medicare policies and coverage, please visit their search website at:

<http://www.cms.gov/medicare-coverage-database/overview-and-quick-search.aspx?from2=search1.asp&> or [the manual website](#)

Clinical Rationale

In 2016, the National Institute for Health and Clinical Excellence (NICE) (NICE, 2016; Singh et al., 2019) issued guidance on therapeutic monitoring of tumor necrosis factor (TNF)-alpha inhibitors in the treatment of patients with Crohn's disease. NICE stated that "enzyme-linked immunosorbent assay (ELISA) kits show promise for therapeutic monitoring of -alpha inhibitors in people with Crohn's disease but there is insufficient evidence to recommend their routine adoption" and recommended that laboratories monitoring (TNF)-alpha inhibitors in patients with Crohn disease who have lost response to the treatment should "work with clinicians to collect data through a prospective study, for local audit, or for submission to an existing registry." NICE also stated that "Enzyme-linked immunosorbent assay (ELISA) tests for therapeutic monitoring of tumor necrosis factor (TNF)-alpha inhibitors (drug serum

levels and antidrug antibodies) show promise but there is currently insufficient evidence to recommend their routine adoption in rheumatoid arthritis.”

In 2017, Feuerstein et al. published the American College of Gastroenterology Institute guidelines on therapeutic drug monitoring in inflammatory bowel disease. The guidelines note that “When anti-drug antibodies are detected, it is unclear what antibody level is clinically meaningful.... The reporting of anti-drug antibodies is variable between commercial assays, with some assays being very sensitive for detecting very-low-titer antibodies of limited clinical significance. Uniform thresholds for clinically relevant antibody titers are lacking. At this time, it is unclear how antibodies affect drug efficacy when both active drug and antibodies are detected. In cases of low trough concentrations and low or high anti-drug antibodies, the evidence to clarify optimal management is lacking.” These guidelines did not address therapeutic drug monitoring in patients treated with vedolizumab or ustekinumab, due to a scarcity of data at the time of publication.

A 2018 retrospective, multicenter study (Papamichael et. al.) reviewed drug retention in inflammatory bowel disease (IBD) patients of whom infliximab was optimized to overcome immunogenicity and variables associated with drug retention. This study consisted of consecutive IBD patients with antibodies to infliximab (ATI), based on either proactive or reactive therapeutic drug monitoring, who underwent infliximab optimization (increasing dose, shortening interval, adding an immunomodulator, or combination) to overcome immunogenicity from September 2012 to July 2015; they were followed through December 2015. ATI were analyzed using the drug-tolerant Prometheus homogeneous mobility shift assay. Drug retention was defined as no need for drug discontinuation due to secondary loss of response (SLR) or serious adverse event. The cohort consisted of 22 patients (Crohn's disease, n = 15). At the end of follow-up [median, (IQR): 17.3 (10.5-32.8) months] 77% (15/22) of patients were still on drug. Univariable Cox proportional hazards regression analysis identified first detectable ATI titer as the only variable associated with drug retention (HR: 0.89; 95% CI: 0.82-0.98, p = 0.016). Receiver-operating characteristic analysis identified an ATI titer < 8.8 U/mL associated with drug retention. The authors concluded that in real-life clinical practice, optimization of infliximab therapy can prevent drug discontinuation in approximately 3/4 of patients with ATI, especially in those with low titers. However, large prospective studies are needed to confirm these data.

A 2019 small cohort study (Van den Berghe et. al.) evaluated ADA to vedolizumab in 40 patients with IBD. This study included the development of an ELISA-based test to measure ADA in the presence of the drug. Anti-vedolizumab antibodies and vedolizumab trough levels were measured after six weeks of treatment and after treatment discontinuation. At the six-week follow-up, three (8%) of the patients were positive for ADA, but this appeared to be transient. None of the patients who discontinued vedolizumab were positive for ADA at the time of their last infusion or after discontinuation. It did not appear that immunogenicity played a major role in vedolizumab treatment failure.

A 2019 clinical utility study (Papamichael et. al.) examined current data regarding therapeutic drug monitoring (TDM) for biologic therapies in IBD. Biologic Therapies include anti-tumor necrosis factor (anti-TNF) agent's infliximab and adalimumab. Up to one-third of patients with Crohn's disease (CD) and ulcerative colitis (UC) show primary non-response (PNR) to biologic therapies, and up to 50% of patients after an initial clinical response stop therapy for either SLR- or a serious adverse event. Both PNR and SLR are due to either pharmacokinetic (PK) or pharmacodynamic (PD) problems. PK issues are associated with inadequate drug exposure, often because of the development of antidrug antibodies (ADA), whereas PD issues are typically related to inflammatory process unrelated to the targeted immunoinflammatory pathway. TDM, defined as the assessment of drug concentrations and ADA, is an important tool for optimizing biologic therapy. However, there are still some limitations when applying TDM into clinical practice, such as when to use TDM, proper interpretation and application of the

results, and the identification of the optimal window/ thresholds to target. These therapeutic windows or thresholds appear to vary on the basis of the outcome of interest and the IBD phenotype. Most of the data on implementation of TDM refer to anti-TNF therapies and the maintenance phase of treatment. The authors concluded that although well-designed large prospective studies are lacking, there are preliminary data mainly from retrospective studies that demonstrate that proactive TDM is associated with better therapeutic outcomes compared with empiric dose optimization and/or reactive TDM. However, major limitations of the evidence relate to the lack of large prospective studies and RCTs on TDM of biologic therapy applied on different IBD phenotypes and sparse data on induction therapy and on biologic agents other than infliximab and adalimumab. Also, it is unclear whether trough concentrations are the best predictor of initial response to biologics, compared with peak drug concentrations or total drug exposure. Further RCTs to establish the utility of proactive TDM, particularly during the induction phase, should be performed. Along with future directions focusing on the development of accurate, easily accessible, and affordable rapid assays, dashboards to allow fast dosing adaptation and incorporation of predictive PK models based on patient and disease characteristics.

In another study published in 2020, Papamichael et. al. assessed the impact of an upward drift for both infliximab and adalimumab concentrations measured by the homogenous mobility shift assay (HMSA) on clinical care of patients with inflammatory bowel disease. Providers reviewed the individual patient data and drug concentrations before and after the laboratory corrections and then documented whether a different clinical decision would have been made had the corrected drug concentration been originally reported. A multivariable Cox proportional hazards regression analysis was performed to investigate the association of a documented treatment change with treatment failure, defined as drug discontinuation for primary nonresponse, loss of response, or serious adverse event, adjusting for confounding factors. The study population consisted of 479 patients (infliximab, n = 219; adalimumab, n = 260). Upon review, 14.9% (71/479) patients would have had a different treatment decision made had the corrected drug concentration been initially reported. After a median follow-up of 10.6 months, 25.7% of patients (123/479) had treatment failure. A theoretical different clinical decision based on the corrected drug concentrations was not associated with treatment failure (adjusted hazard ratio (HR): 1.452; 95% confidence interval (CI): 0.805-2.618; p = 0.216), which was consistent for both infliximab (adjusted HR: 1.977; 95% CI: 0.695-5.627; p = 0.201) and adalimumab (adjusted HR: 1.484; 95% CI: 0.721-3.054; p = 0.284). The authors concluded that the HMSA infliximab and adalimumab assay drift led to providers reporting a change in care of 15% of patients based on the corrected value. However, these decisions did not lead to a higher rate of treatment failure compared to patients without a retrospective (theoretical) treatment change. There were also significant quantitative and qualitative differences between infliximab and adalimumab concentrations before and after the correction of the HMSA, directly influencing drug concentration thresholds associated with clinical outcomes.

Ungaro and associates looked at IBD patients on maintenance vedolizumab, examining the association of maintenance vedolizumab concentrations with remission in their cross-sectional multi-center study from 2019. A homogeneous mobility shift assay was used to determine trough serum concentrations of vedolizumab and anti-drug antibodies (ATVs). The primary outcome was corticosteroid-free clinical and biochemical remission defined as a composite of clinical remission, normalized C-reactive protein [CRP] and no corticosteroid use in 4 weeks. Secondary outcomes included corticosteroid-free endoscopic and deep remission. Vedolizumab concentrations were compared between patients in remission and with active disease. Logistic regression, adjusting for confounders, assessed the association between concentrations and remission. In total, 258 IBD patients were included (55% CD and 45% UC). Patients in clinical and biochemical remission had significantly higher vedolizumab concentrations (12.7 µg/mL vs 10.1 µg/mL, p = 0.002). Concentrations were also higher among patients in endoscopic and deep remission (14.2 µg/mL vs 8.5 µg/mL, p = 0.003 and 14.8 µg/mL vs 10.1 µg/mL, p = 0.01, respectively).

After controlling for potential confounders, IBD patients with vedolizumab concentrations >11.5 µg/mL were nearly 2.4 times more likely to be in corticosteroid-free clinical and biochemical remission. Only 1.6% of patients had ATVs. The authors stated that in a large real-world cohort of vedolizumab maintenance concentrations, IBD patients with remission defined by objective measures (CRP and endoscopy) had significantly higher trough vedolizumab concentrations and immunogenicity was uncommon.

In another study from 2019, a randomized control trial (Hambardzumyan et al.) examined how infliximab-treated patients with rheumatoid arthritis (RA) may respond insufficiently due to low serum infliximab (sIFX) levels, caused by anti-drug antibodies (ADAs). Even though monitoring of sIFX and ADAs is not routinely implemented, and levels for optimal outcome have not been validated. Serum samples of infliximab (sIFX) and anti-drug antibodies (ADAs) levels randomized to methotrexate + infliximab therapy at three, nine, and 21 months were available (n=101). The primary and secondary outcome measures were low disease activity [LDA = 28-joint Disease Activity Score (DAS28) ≤ 3.2] and remission (DAS28 < 2.6). The frequencies of very low sIFX levels increased over time, with 15%, 23%, and 28% at 3, 9, and 21 months from IFX start, respectively, and the majority of patients with very low sIFX levels were ADA positive at these time-points [71% (10/14), 82% (18/22), and 68% (19/28), respectively]. The proportion of patients with LDA was numerically higher at all follow-up time-points among those with sIFX ≥ 0.2 µg/mL compared with patients who had sIFX < 0.2 µg/mL and positive ADAs, although only significant at 21 months (67% and 26%, p=0.002). Similar results were observed when remission was the outcome measure (47% vs 11%, p=0.004). In conclusion, even though monitoring of sIFX and ADAs is not routinely implemented, and levels for optimal outcome have not been validated, these findings support the monitoring of serum drug levels. However, this conclusion requires validation in larger populations and for dose-adjustment studies.

Lastly, in a 2020 review, Papamichael and Cheifetz presented an analysis on the role of TDM of biologics in patients with IBD. Numerous prospective exposure-response relationship studies and post-hoc analyses of RCTs showed a positive correlation between biologic drug concentrations and favorable clinical outcomes in IBD. These studies also demonstrated that higher drug concentrations appeared to be needed to achieve more stringent objective therapeutic outcomes. Reactive TDM rationalizes the management of primary non-response and secondary loss of response to anti-TNF therapy and is more cost-effective when compared with empiric dose optimization. Additionally, recent data suggest that proactive TDM, with the goal of targeting a threshold drug concentration, is associated with better therapeutic outcomes when compared with empiric dose escalation and/or reactive TDM of infliximab or adalimumab. Finally, proactive TDM can also efficiently guide infliximab de-escalation or discontinuation in patients with IBD in remission. The authors found that reactive TDM is currently considered as standard of care, whereas proactive TDM is emerging as a new therapeutic strategy for better optimizing anti-TNF therapy in IBD. Nevertheless, further data from prospective studies are needed before a wide implementation of TDM-based algorithms in real life clinical practice for newer biologics.

Applicable Coding

CPT Codes

80145	Adalimumab
80230	Infliximab
80280	Vedolizumab

80299 Quantitation of therapeutic drug, not elsewhere specified

84999 Unlisted chemistry procedure

HCPCS Codes

Not Applicable

References:

1. Centers for Medicare and Medicaid Services (CMS). CMS Manual System. Pub 100-04 –Medicare Claims Processing. Transmittal 2639. Date: Jan. 25, 2013. SUBJECT: Healthcare Common Procedure Coding System (HCPCS) Codes Subject to and Excluded from Clinical Laboratory Improvement Amendments (CLIA) Edits. See <http://www.cms.hhs.gov/CLIA/>.
2. Feuerstein JD, Nguyen GC, Kupfer SS, et al. American Gastroenterological Association Institute Guideline on Therapeutic Drug Monitoring in Inflammatory Bowel Disease. *Gastroenterology*. Sep 2017; 153(3): 827-834. PMID 28780013
3. Hambardzumyan, K, Hermanrud, C, Marits, P, et al. Association of female sex and positive rheumatoid factor with low serum infliximab and anti-drug antibodies, related to treatment failure in early rheumatoid arthritis: results from the SWEFOT trial population. *Scandinavian journal of rheumatology*. 2019 Sep;48(5):362-6. PMID: 31244356
4. National Institute for Health and Care Excellence (NICE).NICE. (2019). "Therapeutic monitoring of TNF-alpha inhibitors in rheumatoid arthritis". Accessed: June 24, 2021. Available at: <https://www.nice.org.uk/guidance/dg36/chapter/1-Recommendations>
5. National Institute for Health and Care Excellence (NICE). Therapeutic monitoring of TNF-alpha inhibitors in Crohn's disease (LISA-TRACKER ELISA kits, IDKmonitor ELISA kits, and Promonitor ELISA kits) [DG22]. 2016; Accessed JuneOctober 2413, 20210. Available at: <https://www.nice.org.uk/guidance/dg22/chapter/1-Recommendations>
6. Papamichael, K. and A. S. Cheifetz (2020). "Therapeutic drug monitoring in patients on biologics: lessons from gastroenterology." *Curr Opin Rheumatol* 32(4): 371-379.
7. Papamichael, K., et al. (2018). "Long-Term Outcome of Infliximab Optimization for Overcoming Immunogenicity in Patients with Inflammatory Bowel Disease." *Dig Dis Sci* 63(3): 761-767.
8. Papamichael, K., et al. (2019). "Appropriate Therapeutic Drug Monitoring of Biologic Agents for Patients with Inflammatory Bowel Diseases." *Clin Gastroenterol Hepatol* 17(9): 1655-1668 e1653.
9. Papamichael, K., et al. (2020). "Clinical Impact of Corrections to Infliximab and Adalimumab Monitoring Results with the Homogeneous Mobility Shift Assay." *J Clin Med* 9(9).
10. Promethius Anser. Integrated IBD Monitoring. Available at: <https://www.anserifx.com/>
11. Singh, J. A., Guyatt, G., Ogdie, A., Gladman, D. D., Deal, C., Deodhar, A., Reston, J. (2019). 2018 American College of Rheumatology/National Psoriasis Foundation Guideline for the Treatment of Psoriatic Arthritis. *Arthritis Care & Research*, 71(1), 2-29. doi:10.1002/acr.23789
12. Ungaro RC, Yarur A, Jossen J, et al. Higher trough vedolizumab concentrations during maintenance therapy are associated with corticosteroid-free remission in inflammatory bowel disease. *J Crohns Colitis*. 2019;13(8):963-969
13. Van den Berghe, N, Verstockt, B, Tops, S, Ferrante, M, Vermeire, S, Gils, A. Immunogenicity is not the driving force of treatment failure in vedolizumab-treated inflammatory bowel disease patients. *Journal of gastroenterology and hepatology*. 2019 Jul;34(7):1175-81. PMID: 30589948

Disclaimer:

This document is for informational purposes only and should not be relied on in the diagnosis and care of individual patients. Medical and Coding/Reimbursement policies do not constitute medical advice, plan preauthorization, certification, an explanation of benefits, or a contract. Members should consult with appropriate health care providers to obtain needed medical advice, care, and treatment. Benefits and eligibility are determined before medical guidelines and payment guidelines are applied. Benefits are determined by the member's individual benefit plan that is in effect at the time services are rendered.

The codes for treatments and procedures applicable to this policy are included for informational purposes. Inclusion or exclusion of a procedure, diagnosis or device code(s) does not constitute or imply member coverage or provider reimbursement policy. Please refer to the member's contract benefits in effect at the time of service to determine coverage or non-coverage of these services as it applies to an individual member.

U of U Health Plans makes no representations and accepts no liability with respect to the content of any external information cited or relied upon in this policy. U of U Health Plans updates its Coverage Policies regularly, and reserves the right to amend these policies and give notice in accordance with State and Federal requirements.

No part of this publication may be reproduced, stored in a retrieval system or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from U of U Health Plans.

"University of Utah Health Plans" and its accompanying logo, and its accompanying marks are protected and registered trademarks of the provider of this Service and or University of Utah Health. Also, the content of this Service is proprietary and is protected by copyright. You may access the copyrighted content of this Service only for purposes set forth in these Conditions of Use.

© CPT Only – American Medical Association